Saskatchewan
Ministry of
Agriculture

Crop Report

Published by the Ministry of Agriculture
ISSN 0701 7085
Report number 12, June 24, 2010

Crop Report for the Period June 15 to 21, 2010

Saskatchewan farmers have seeded 76 per cent of the 2010 crop, according to Saskatchewan Agriculture's weekly Crop Report. In general, the past weeks' rain has made seeding stressful and left many acres unseeded. It has caused flooding in many areas. Thunderstorms continue to roll through the province.

Seeding is 96 per cent complete in northwestern Saskatchewan; 89 per cent complete in southwestern Saskatchewan; 91 per cent complete in west-central Saskatchewan; 76 per cent complete in southeastern Saskatchewan; 59 per cent complete in east-central Saskatchewan and 50 per cent complete in the northeast.

Across the province, cropland topsoil moisture is rated as 64 per cent surplus and 36 per cent adequate. Topsoil moisture on hay land and pasture is rated as 39 per cent surplus, 60 per cent adequate and one per cent short.

Fifty-two per cent of the province's fall-seeded cereals are at a normal stage of development, but 77 per cent of the spring-seeded cereals, 77 per cent of the oilseeds and 67 per cent of the pulses are behind normal in development.

Seventy-five and 80 per cent of the winter wheat and fall rye are in good to excellent condition, respectively. Seventy-seven per cent of the spring wheat, 84 per cent of the durum, 70 per cent of the barley, 72 per cent of the oats, 75 per cent of the flax, 71 per cent of the canola, 79 per cent of the mustard, 83 per cent of the lentils, 82 per cent of the peas, 78 per cent of the canaryseed and chickpeas are in good to fair condition.

Crops are showing signs of stress from excess moisture. Crop damage is attributed mostly to flooding and hail. Cutworm, wireworm and gopher damage has been reported, along with leaf diseases and root rots.

Haying has yet to begin for the majority of producers. Farmers are controlling weeds.

Southeastern Saskatchewan (Crop Districts 1, 2, and 3ASE)

The region has approximately 76 per cent of the 2010 crop in the ground. The region received an average of 54 mm of moisture during the past week. The Manor area received 92 mm, the Tantallon area 76 mm, the Weyburn area 63 mm, the Marquis area 132 mm and the Ceylon area 36 mm. So far this year, the region has received between 158 and 350 mm of rain.

Topsoil moisture conditions on cropland are reported as 73 per cent surplus and 27 per cent adequate. On hay land and pasture, topsoil moisture is rated as 42 per cent surplus, 57 per cent adequate and one per cent short.

One year ago
Cutworms were causing damage in the northeastern regions. Majority of crop damage was attributed to dry conditions.

For further information, contact: Grant McLean, PAg,
Cropping Management Specialist, Agriculture Knowledge Centre,
Toll Free: 1-866-457-2377, E-mail: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at www.agriculture.gov.sk.ca.

Agriculture and Agri-Food Canada Agriculture et Agroalimentaire Canada

CANADA – SASKATCHEWAN
Crop Insurance

Crop damage is mostly attributed to flooding. Flea beetle and leaf diseases are causing damage as well. Hail was reported in the Odessa area.

SE SK Crop Development – June 21, 2010	% Ahead	% Normal	% Behind
Fall Cereals	7	82	11
Spring Cereals	0	28	72
Oilseeds	0	30	70
Pulse Crops	1	48	51

Warm and sunny weather is needed as crops in areas are under moisture stress. Weeds are becoming an issue. Farmers are busy with weed control as weeds are starting to get ahead of some crops. Farmers are applying fungicides. Wet areas in the fields are making navigation difficult at best. Areas north and west of Carlyle were flooded after five to seven inches of rain fell on June 18. Farmers are trying to assess the extent of flooding damage to crops. Crop development is variable because some crops were seeded early while others were just seeded late last week. Producers are trying roll pulses when the weather co-operates. Some farmers are planning to seed greenfeed or to plough down crops to use up some of the excess moisture in the fields. Hay and pasture continue to look good.

Southeast SK Crop conditions- June 21, 2010	Winter Wheat	Fall Rye	Spring Wheat	Durum	Oat	Barley	Flax
% excellent	24	62	9	8	6	7	2
% good	63	22	39	30	41	34	35
% fair	12	14	35	44	37	37	42
% poor	1	0	14	15	14	19	18
% very poor	0	2	3	3	2	3	3
	Canola	Mustard	Pea	Lentil	Canary-seed	Chick-pea	
% excellent	5	0	6	4	0	0	
% good	35	34	44	37	20	33	
% fair	40	44	33	40	60	54	
% poor	18	21	15	17	19	7	
% very poor	2	1	2	2	1	6	

Southwestern Saskatchewan (Crop Districts 3ASW, 3AN, 3B and 4)

Eighty-nine per cent of the 2010 crop has been seeded. The acres that are left unseeded are still too wet to even attempt seeding. Crop reporters indicated some farmers are trying to seed annual forage crops. Parts of the region received significant rainfall last week, resulting in flooding and damage to roads, highways, fields and yards. The Maple Creek area received the worst of the downpour – over seven inches of rain fell overnight on June 18. The region received an average of 62 mm of rain. The Limerick area received 30 mm, the Chaplin area 63 mm, the Shaunavon area 85 mm, the Lucky Lake area 81 mm, the Gull Lake area 173 mm, the Big Beaver area 14 mm and the Bengough area 19 mm. Other areas receiving more than 100 mm of rain for the week include Richmound, Fox Valley and Leader. So far for this year, the region has received between 139 and 376 mm of rain.

Cropland topsoil moisture is rated as 47 per cent surplus and 53 per cent adequate. Hay land and pasture topsoil moisture conditions are 24 per cent surplus, 75 per cent adequate and one per cent short. Crop District 3AN is reporting 83 per cent of the cropland has surplus topsoil moisture.

Crop damage has been caused by flooding and hail. Hail damage was reported in the Spring Valley, Mossbank, Glenbain, Blumenhof, Shaunavon and Consul areas. Cutworms and wireworms are causing damage in the Webb area. Gophers are also causing crop damage.

SW SK Crop Development – June 21, 2010	% Ahead	% Normal	% Behind
Fall Cereals	1	41	58
Spring Cereals	0	17	83
Oilseeds	0	17	83
Pulse Crops	0	21	79

Sun and heat are needed to relieve the seeded crops of excess moisture stress. Some areas are reporting that crops are looking good and emergence is even. Ascochyta is starting to show up in chickpeas. A tornado touched down in the Limerick area on June 21, damaging bins and sheds. Farmers are busy trying to control weeds and applying top-up fertilizer where crops are showing signs of nutrient deficiency. Some farmers are rolling pulses despite the fact that their crops are too far advanced for safe rolling. Crop staging is variable across the region. The hay crop looks great. Cattle are doing well on pasture.

Southwest SK Crop Conditions- June 21, 2010

	Winter Wheat	Fall Rye	Spring Wheat	Durum	Oat	Barley	Triticale
% excellent	25	27	12	11	8	12	25
% good	49	58	50	48	54	55	47
% fair	23	14	37	39	37	32	28
% poor	3	1	1	2	1	1	0
% very poor	0	0	0	0	0	0	0

	Flax	Canola	Mustard	Pea	Lentil	Canary-seed	Chick-pea
% excellent	10	8	23	13	11	7	9
% good	37	59	60	49	37	65	52
% fair	52	31	17	37	49	28	24
% poor	1	2	0	1	3	0	15
% very poor	0	0	0	0	0	0	0

East-Central Saskatchewan (Crop Districts 5 and 6a)

Farmers in the region have 59 per cent of the crop in the ground. Heavy rain throughout the week did not help farmers deal with flooded acres or access land with equipment. The region received an average of 33 mm of rain. Seeding in the east-central region continues to be impeded by saturated soil and continuous rain. The Neudorf area recorded 77 mm of rain, the Foam Lake area received 50 mm, the Jansen area 52 mm and the Lintlaw and Raymore areas received 17 mm. So far for this year, the region has received between 158 and 404 mm of rain.

Topsoil moisture continues to increase as the rain continues. Cropland topsoil moisture is reported as 88 per cent surplus and 12 per cent adequate. Hay land and pasture topsoil moisture is rated as 69 per cent surplus and 31 per cent adequate.

Flooding is causing the majority of crop damage. Cutworms, flea beetles and leaf diseases are also damaging crops.

Low-lying areas continue to be flooded. Some farmers are still trying to seed. The recent brief break in wet weather in some areas provided some relief to crops that were water-stressed and behind normal in development. Farmers are controlling weeds and dodging sloughs. Weeds are too-far-advanced in growth to be easily controlled. Emergence is uneven is some areas because the seed rotted in the soil before it had a chance to emerge. Hay fields and pastures were doing well, but are also flooded in areas. Warm and dry weather is needed before haying equipment can get into the forage fields. Parts of some pastures cannot be grazed because of flooding. One crop reporter indicated this is the wettest spring on his farm since the family started keeping records in 1948.

EC SK Crop Development
June 21, 2010

	% Ahead	% Normal	% Behind
Fall Cereals	0	48	52
Spring Cereals	0	15	85
Oilseeds	4	23	73
Pulse Crops	0	30	70

East -Central SK Crop Conditions- June 21, 2010

	Winter Wheat	Fall Rye	Spring Wheat	Durum	Oat	Barley
% excellent	25	30	2	5	2	4
% good	45	40	34	50	35	36
% fair	17	19	31	37	25	29
% poor	13	11	31	8	38	25
% very poor	0	0	2	0	0	6

	Flax	Canola	Mustard	Pea	Lentil	Canary-seed
% excellent	4	4	6	9	9	9
% good	40	32	40	42	51	12
% fair	29	28	30	30	20	31
% poor	22	32	24	18	20	48
% very poor	5	4	0	1	0	0

West-Central Saskatchewan (Crop Districts 6B and 7)

Ninety-one per cent of the region's crops have been seeded. An average of 56 mm of rain fell on the region. The Hanley area received 59 mm, the Eston area 129 mm, the Kindersley area 123 mm, the Brock area 107 mm and the Biggar area 65 mm. The Sonningdale and Macklin areas recorded the least amount of precipitation – 24 mm. So far for this year, the region has received between 188 and 320 mm of rain.

WC SK Crop Development – June 21, 2010

	% Ahead	% Normal	% Behind
Fall Cereals	8	54	38
Spring Cereals	2	26	72
Oilseeds	4	23	73
Pulse Crops	2	33	65

Cropland topsoil moisture is rated as 53 per cent surplus and 47 per cent adequate. Hay land and pasture topsoil moisture conditions are rated as 34 per cent surplus and 66 per cent adequate.

The majority of crop damage is due to flooding. Cutworm and leaf disease damage is showing up. Hail damage was reported in the Kerrobert area.

The week's rain stopped seeding and weed control operations. Weed control is an issue as some fields are very wet and some crops are getting past the weed control stage. Some farmers will be putting in oats or barley for greenfeed. Flooding has resulted in some crop loss. Crops are behind normal in development, but look to be in good condition, especially where there is good drainage. However, entire crops have turned yellow in areas. Crops progressed quickly during the few days of warm, sunny weather. Hay and pasture are looking very good.

West-Central SK Crop Conditions- June 21, 2010

	Winter Wheat	Fall Rye	Spring Wheat	Durum	Oat	Barley	Flax
% excellent	7	5	1	6	2	1	0
% good	75	79	65	67	65	19	64
% fair	9	14	27	23	31	24	29
% poor	9	2	7	3	2	48	7
% very poor	0	0	0	1	0	8	0

	Canola	Mustard	Pea	Lentil	Canary-seed	Chick-Pea
% excellent	2	6	7	9	9	1
% good	60	70	70	70	67	53
% fair	28	23	21	19	21	44
% poor	9	1	2	1	2	2
% very poor	1	0	0	0	1	0

Northeastern Saskatchewan (Crop Districts 8 and 9AE)

The northeastern region has 50 per cent of the 2010 crop in the ground. No seeding progress was made during the week. An average of 37 mm of rain fell on already saturated fields. The Tisdale area received 49 mm, the Melfort area 50 mm and the Garrick area 31 mm. So far this year, the region has received between 163 and 332 mm of rain.

Cropland topsoil moisture is rated as 89 per cent surplus and 11 per cent adequate. Hay land and pasture topsoil moisture is rated as 81 per cent surplus and 19 per cent adequate.

NE SK Crop Development
June 21, 2010

	% Ahead	% Normal	% Behind
Fall Cereals	0	51	49
Spring Cereals	0	13	87
Oilseeds	0	12	88
Pulse Crops	0	18	82

Crop damage is due to flooding. Cutworms are causing damage in the Star City, Arborfield and Nipawin areas. Leaf diseases and root rots are also causing crop damage. Hail was reported in the Garrick area, and the extent of the damage is still being assessed.

Crop reporters are indicating that seeding is done for 2010. Farmers are now concentrating on weed control and dealing with excess moisture in fields. Weeds are starting to get ahead of crops. Entire crops have turned yellow in some areas. Crop conditions and development are variable due to the extended and wet seeding season. The couple of days of sunny weather during the week provided some relief for stressed crops.

Northeast SK Crop Conditions- June 21, 2010					
	Winter Wheat	Fall Rye	Spring Wheat	Oat	Barley
% excellent	0	0	1	1	8
% good	29	41	19	9	63
% fair	16	24	29	35	29
% poor	54	34	44	52	0
% very poor	1	1	7	3	0
	Flax	Canola	Mustard	Pea	Lentil
% excellent	0	1	0	1	0
% good	6	22	20	24	50
% fair	52	34	40	32	35
% poor	36	34	25	40	10
% very poor	6	9	0	3	5

Northwestern Saskatchewan (Crop Districts 9AW and 9B)

Seeding is 96 per cent complete for the region. The rains missed much of the region, allowing producers to get some field work done. The region received an average of 7 mm of rain. The Hafford area received 32 mm; the North Battleford area 20 mm. Many areas received less than 10 mm of moisture. The Turtleford and Frenchman Butte areas recorded no rain for the week. So far for this year, the region has received between 146 and 291 mm of rain.

NW SK Crop Development – June 21, 2010			
	% Ahead	% Normal	% Behind
Fall Cereals	0	43	57
Spring Cereals	8	40	52
Oilseeds	2	43	55
Pulse Crops	6	52	42

Topsoil moisture conditions for cropland are four per cent surplus, 92 per cent adequate and four per cent short. Pasture and hay land conditions are two per cent surplus, 95 per cent adequate and three per cent short.

The majority of crop damage was caused by flooding. Cutworm damage was reported in the Speers, North Battleford and Barthel areas.

Crops are emerging and looking good for the most part, but are behind in development. Farmers are putting in greenfeed and crops for swath-grazing. Some low-lying acres have been lost to standing water. Early seeded crops are looking better than the later seeded crops. Farmers are controlling weeds. Some have indicated a second herbicide application is warranted due to the later flush of weeds. Applications of cutworm control were reported in the Speers area. Heat and sun would be appreciated to get things going. Some crops are yellowing under the heavy rain due to excess moisture stress. Hay and pastures look good.

North west SK Crop Conditions-June 21, 2010

	Winter Wheat	Fall Rye	Spring Wheat	Oat	Barley
% excellent	0	50	1	2	17
% good	100	40	68	74	51
% fair	0	10	30	23	25
% poor	0	0	1	1	6
% very poor	0	0	0	0	1

	Flax	Canola	Mustard	Pea	Lentil
% excellent	0	2	N/A	1	N/A
% good	100	68	N/A	71	N/A
% fair	0	29	N/A	27	N/A
% poor	0	1	N/A	1	N/A
% very poor	0	0	N/A	0	N/A

Crop Districts and Rural Municipalities in Saskatchewan



Weekly Rainfall Summary

(in millimeters) 1 inch = 25 mm for the period June 15 to 21, 2010

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
1A	1	Argyle	N/A	205.0
	2	Mount Pleasant	70.0	206.0
	3	Enniskillen	89.0	226.0
	33 A	Moose Creek	N/A	39
	33 B	Moose Creek	N/A	233.0
	34	Browning	29.0	208.0
	61	Antler	N/A	186.0
	63	Moose Mountain	92.0	249.0
	64	Brock	37.0	206.0
	65	Tecumseh	40.0	239.0
1B	91	Maryfield	35.0	223.0
	92	Walpole	58.0	216.0
	93	Wawken	50	232
	95	Golden West	64	263
	122	Martin	67	350.0
	123 A	Silverwood	52	204
	123 B	Silverwood	N/A	N/A
	124	Kingsley	36	183
	125	Chester	45	214
	151 A	Rocanville	76	281
	151 B	Rocanville	58	296
	153	Willowdale	54	262
	154	Elcapo	28	243
	155	Wolseley	53	286
2A	66	Griffin	N/A	222
	67	Weyburn	63	245
	68	Brokenshell	45	222
	69	Norton	29	283
	96	Fillmore	43	204
	97	Wellington	36	158
2B	127 A	Francis	59.5	235
	127 B	Francis	74	180
	129	Bratt's Lake	42	172
	156 A	Indian Head	50.5	252
	156 B	Indian Head	67	250
	157	South Qu'appelle	N/A	NIL
	160	Pense	47	213
	161	Moose Jaw	60	253
	191	Marquis	132	333
3ASE	38 A	Laurier	30	226
	38 B	Laurier	22	201
	39 A	The Gap	35	228
	39 B	The Gap	36	297
3ASW	10	Happy Valley	14	307
	11	Hart Butte	N/A	NIL
	12	Poplar Valley	N/A	198
	40	Bengough	19	256
	42	Willow Bunch	20	287
	43	Old Post	23	211
	44	Waverley	N/A	203
	70	Key West	N/A	174
	71 A	Excel	N/A	192
	71 B	Excel	14	198
	73 A	Stonehenge	27.7	264
	73 B	Stonehenge	30	326
	74	Wood River	23	180
3AN	101	Terrell	53	306
	102	Lake Johnston	29.5	185
	103	Sutton	46	251
	134	Shamrock	54	291
	164	Chaplin	63	271
	193	Eyebrow	58	279
	194	Enfield	N/A	241
3BS	17	Val Marie	27	195
	45	Mankota	33	215
	75 A	Pinto Creek	40	210
	75 B	Pinto Creek	27	225
	76 A	Auvergne	13	164
	76 B	Auvergne	20	155
	77	Wise Creek	29	176
	78 A	Grassy Creek	N/A	192
	78 B	Grassy Creek	75	244
	78 C	Grassy Creek	85	287
	105	Glenbain	25	189
	106	Whiska Creek	21	139
	107	Lac Pelletier	52	201
	108	Bone Creek	70	252
3BN	138	Webb	72.5	267
	165	Morse	N/A	NIL
	166	Excelsior	59	367
	167	Sask. Landing	47.78	266
	168	Riverside	80	280
	226	Victory	38	276
	228	Lacadena	81	269
	255	Coteau	N/A	NIL
	257	Monet	60	229

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
4A	49	White Valley	68	246
	49 A	White Valley	58	198
	51	Reno	85.4	224
	79	Arlington	133	293
	109	Carmichael	173	376
	110	Piapot	125	306
	111	Maple Creek	122	328
4B	139	Gull Lake	158	349
	141	Big Stick	107	332
	142	Enterprise	121	355
	169	Pittville	83	286
	171	Fox Valley	119	287
	229	Miry Creek	84	318
	231	Happyland	124	258
5A	181	Langenherg	30	230
	183	Fertile Belt	48	234
	185	Mcleod	77	241
	186	Abernethy	33	207
	211	Churchbridge	26	253
	213	Saltcoats	16	230
	216	Tullymet	N/A	147
	241	Calder	60	265
	243	Wallace	24	235
	245 A	Garry	32	312
	245 B	Garry	38	330
	246	Ituna Bon Accord	22	198
	247	Kellross	35	260
	248	Touchwood	17	158
5B	271	Cote	18	217
	275	Insinger	N/A	NIL
	276	Foam Lake	50	360
	277	Emerald	32	335
	278	Kutawa	N/A	NIL
	305	Invermay	34	361
	307	Elfros	40	404
	308	Big Quill	30	271
	333	Clayton	N/A	133
	335	Hazel Dell	17	250
	336	Sasman	24	293
	337	Lakeview	N/A	272
	338	Lakeside	46	320
	366	Kelvington	24	237
6A	190 A	Dufferin	25	174
	190 B	Dufferin	43	255
	190 C	Dufferin	51	209
	190 D	Dufferin	1.5	272
	219 A	Longlaketon	24	182
	219 B	Longlaketon	28	199
	220	Mckillop	27	208
	221	Sarnia	46	266
	222	Craik	26	266
	251	Big Arm	31	273
	252	Arm River	36	247
	279	Mount Hope	18	256
	280	Wreford	25	278
	282	McCraney	26	306
	309	Prairie Rose	52	345
	313	Lost River	32	249
	339	Leroy	46.4	340
	340	Wolverine	N/A	295
	341	Viscount	45	324
	343	Blucher	37	292
6B	253	Willner	30	221
	283	Rosedale	25	305
	284	Rudy	N/A	309
	285	Fertile Valley	37	282
	286	Milden	48	291
	314	Dundurn	59	293
	344	Corman Park	44	285
	346	Perdue	N/A	290
	376 A	Eagle Creek	79	398
	376 B	Eagle Creek	24	284
7A	260	Newcombe	129	259
	260 B	Newcombe	123	188
	287	St. Andrews	56	233
	288	Pleasant Valley	61	247
	290 A	Kindersley	87	196
	290 B	Kindersley	107	247
	292	Milton	68	205
	317 A	Marriott	54	285
	317 B	Marriott	68	320
	318 A	Mountain View	69	242
	318 B	Mountain View	59	291
	320	Oakdale	78	242
	321	Prairiedale	90	264

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
7B	347 A	Biggar	38	246
	347 B	Biggar	65	269
	350 A	Manposa	56	209
	350 B	Manposa	46	200
	351	Progress	38	247
	352	Heart's Hill	56	260
	377	Glenside	25	315
	378 A	Rosemount	N/A	157
	378 B	Rosemount	52	288
	379	Reford	45	252
	380	Tramping Lake	30	209
	381	Grass Lake	50	294
	382	Eye Hill	N/A	206
	409	Buffalo	40	320
	410	Round Valley	10	204
	411	Senlac	24	210
8A	395	Porcupine	22	235
	397	Barrier Valley	38.8	287
	428	Star City	44	246
	456	Arborfield	32	194
	457	Connaught	49	305
	458	Willow Creek	N/A	210
	486	Moose Range	33	206
8B	369	St. Peter	45	303
	370	Humboldt	30	362
	371	Bayne	48	329
	372	Grant	27	286
	402	Fish Creek	28	249
	429 A	Flett's Springs	38	163
	429 B	Flett's Springs	50	242
	430	Invergordon	N/A	43
	459	Kinistino	N/A	246
9AE	461	Prince Albert	N/A	263
	488	Torch River	31	315
	520	Paddockwood	N/A	67
9AW	406	Mayfield	17	211
	435	Redberry	32	291
	436	Douglas	6	159
	463	Duck Lake	N/A	190
	464	Leask	N/A	265
	467	Round Hill	N/A	NIL
	494	Canwood	N/A	229
	496	Spiritwood	13	276
	497	Medstead	4	186
9B	438	Battle River	20	262
	440	Hillsdale	4	204
	442	Manitou Lake	8.2	203
	498 A	Parkdale	2	228
	498 B	Parkdale	N/A	113
	499	Mervin	NIL	174
	501 A	Frenchman Butte	4	146
	501 B	Frenchman Butte	NIL	188
	501 C	Frenchman Butte	NIL	188
	502	Britannia	3	201
	561	Loon Lake	2	187
	588 A	Meadow Lake	14	223
	588 B	Meadow Lake	N/A	209
	588 C	Meadow Lake	2	227
	588 D	Meadow Lake	N/A	182
	622	Beaver River	2	223

Municipality No: A, B, C, and D - more than one reporter

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.

Weekly Rainfall

for the week ending June 21, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: June 23, 2010

Cumulative Rainfall

From: April 1, 2010
To: June 21, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: June 23, 2010

Weekly Rainfall

for the week ending June 21, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

0 25 50 100 150 200
Kilometers

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: June 23, 2010

Cumulative Rainfall

From: April 1, 2010
To: June 21, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

0 25 50 100 150 200
Kilometers

N

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: June 23, 2010

Cropland Topsoil Moisture Conditions

June 22, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: June 23, 2010

Hay and Pasture Topsoil Moisture Conditions

June 22, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: June 23, 2010

Cropland Topsoil Moisture Conditions

June 22, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

©2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

Projection: UTM Zone 13, Datum: NAD83

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: June 23, 2010

Hay and Pasture Topsoil Moisture Conditions

June 22, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: June 23, 2010